

SOCIETY OF PHARMACEUTICAL EDUCATION & RESEARCH [SPER]

In Collaboration with **VYWS'S INSTITUTE OF PHARMACEUTICAL EDUCATION AND RESEARCH**



Annual International Conference & Exhibition (SPER 2019)



February 22-23, 2019

Jointly Organized by: VYWS'S Institute Of Diploma In Pharmacy (Govt. Aided)

Venue: VYWS'S Institute of Pharmaceutical Education and Research Borgaon Meghe, Wardha [Maharashtra] India

Knowledge Partner

SPER Official Publication

Journal of Advanced Pharmaceutical Technology & Research [JAPTR]



IFTM University, Moradabad [Uttar Pradesh]

Publication Partner



Supported By

Industry Partner

Academic Partner

Media Partner











SPER Times [A Pharma Magazine Conjoining Corporate with Academial



PH-01

EFFECT OF CALORIC VESTIBULAR STIMULATION ON BRAIN NEUROTRANSMITTERS IN A MPTP-INDUCED MOUSE MODEL OF PARKINSON'S DISEASE

Archana R^{1*}, Mukkadan J K², G. Parvathi Mani Pravallika³, Kumar Sai Sailesh⁴

¹Department of Physiology, Saveetha Medical College, Thandalam, Chennai, Tamil Nadu

²Little Flower Medical Research Centre, Angamaly, Kerala, India

³Vishnu Dental College, Vishnupur, Bhimavaram, West Godavari District, Andhra Pradesh

⁴Department of Physiology, Vishnu Dental College, Vishnupur, Bhimavaram

ABSTRACT

The Parkinson disease is a slow progressive, degenerative disease of nervous system associated with damage of dopaminergic neurons in the brain. Though PD was mainly due to damage of damage of dopaminergic system, other neuro-transmitters also effected during PD and contribute to non-motor symptoms of PD. The current study was undertaken to observe the effect of caloric vestibular stimulation on brain neurotransmitters in a MPTP-induced mouse model of Parkinson's disease. 24 healthy, adult male Swiss albino mice with body weight ranging between 25 - 40g were used in the study. The middle ear cavity of the mice was irrigated with hot (40°C) water. 0.5 ml of water was taken in 5 ml syringe with the needle removed. The ear was irrigated with water drop by drop, using the syringe. After 30 days of experimental period, the animals were fasted overnight and sacrificed by cervical decapitation and neuro-transmitter levels were estimated. The present study provides evidence for beneficial effects of caloric vestibular stimulation in limiting the changes in neuro-transmitter levels in Parkinson's disease. We recommend further detailed studies in this area to understand the mechanism of action and to recommend vestibular stimulation as an adjunctive therapy in the management of Parkinson's disease.

PH-02 DEVELOPMENT OF PICTOGRAMS FOR PATIENT EDUCATION Gunja Joshi*, Ajay Pise

Dadasaheb Balpande College of Pharmacy, Nagpur, Maharashtra, India - 440037

ABSTRACT

Aim was to understand and analyse concept of pictogram in educating patients and to develop Pictograms for Patient Education. One of the sources of poor health outcomes is the lack of compliance with the prescribed treatment plans, often due to communication barriers between healthcare professionals and patients. Pictograms are a form of communication that conveys meaning through its pictorial resemblance to a physical object or an action. Pharmaceutical pictograms are often associated with a better comprehension of treatment regimens, although their use is still subject to limitations. This study aims to understand and analyse concept of pictogram in educating patients and to develop pictograms for patient education. Methodology includes collecting information on use of pictograms in patient education through primary and secondary data collection methods, analysis of collected information to understand role of pictogram in patient education and development of designs of pictograms and its evaluation.

PH-03

EFFECT OF CALORIC VESTIBULAR STIMULATION ON BRAIN NEUROTRANSMITTERS IN A ROTENONE-INDUCED MOUSE MODEL OF PARKINSON'S DISEASE

Baavana Bantupalli*, Kumar Sai Sailesh

Vishnu Dental College, Bhimavaram, West Godavari District, Andhra Pradesh

ABSTRACT

The Parkinson disease is a slow progressive, degenerative disease of nervous system associated with damage of dopaminergic neurons in the brain. Though PD was mainly due to damage of dopaminergic system, other neurotransmitters also effected during PD and contribute to non-motor symptoms of PD. The current study was